

PETRYK M.R.¹, BOYKO I.V.¹, SHYNKARYK M.I.², PETRYK O.JU.¹

NONLINEAR MODEL OF THE THREE-COMPONENTS COMPETITIVE ADSORPTION USING LANGMUIR EQUILIBRIUM

The basis for the mathematical modeling of non-isothermal gas competitive adsorption in a porous solid using Langmuir equilibrium is given. High-performance analytical solutions of considered adsorption models based on the Heaviside operating method and Landau's decomposition and linearization approach of Langmuir equilibrium by expanding into a convergent series in the temperature phase transition point are proposed.

The numerical experiments result based on high-speed computations on multicore computers are presented.

Key words and phrases: competitive adsorption, nanoporous medium, diffusion coefficient, competitive Langmuir equilibrium, Heaviside operating method, high-performance calculations.

¹Ternopil Ivan Puluji National Technical University, 56 Ruska str., 46001 Ternopil, Ukraine

²West Ukrainian National University, 11 Lvivska str., 46009 Ternopil, Ukraine

e-mail: *Mykhaylo_Petryk@tu.edu.te.ua*, *jacques.fraissard@upmc.fr*

INTRODUCTION

The experimental and theoretical study of the competitive adsorption and diffusion of several gases through a nanoporous solid and the instantaneous (out of equilibrium) distribution of the adsorbed phases is particularly important in many fields, such as gas separation, heterogeneous catalysis, purification of confined atmospheres, reduction of exhaust emissions contributing to global warming, etc. [1]. Taking into account the influence of physical factors that limit the adsorption kinetics on the surface of nanopores, the quality of the mathematical models for the adsorption of exhaust gases (hydrocarbon components, CO₂) in a microporous bed determines the effectiveness of technological solutions for the neutralization of gas emissions [2–8].

However, most of these models do not fully reflect the complex spatial-temporal representations of the course of all components of complex mass transfer in the intercrystallite space and in the intracrystallite space, including the internal kinetics of the phase transition taking into account the geometric characteristics of transfer areas [6, 7].

In proposed paper, which is the development of papers [8–11], substantiated and developed highly productive methods for mathematical modeling of three-component adsorption

UDC 517.444

2010 *Mathematics Subject Classification:* 46F10, 65R10.

in the microporous solid based on a system of spatiotemporal equations of heat and mass transfer in partial derivatives and generalized Langmuir equation. For modeling, we use the high-performance methods of the Heaviside operational calculus and the decomposition approach for expansion the adsorption equilibrium.

1 COMPETITIVE N -COMPONENT ADSORPTION MODEL IN GENERAL FORMULATION

The presented model is analogous to the biporous model [2, 3, 5, 6]. Developing the approach described by Rhutwen and Karger [7, 8] and Petryk et al. [9] concerning the construction of a complex process of competitive adsorption and diffusion, one should dwell on the most important defining hypotheses limiting the process.

The general hypothesis adopted in developing the presented model in the most general formulation concludes that the competitive n -component adsorption interaction between adsorption molecules of several gases (two or more) and active adsorption centers on the phase separation surfaces in the nanoporous crystallites is determined on the basis of the nonlinear competitive equilibrium function of the Langmuir type, taking into account the physical assumptions [7]:

1. Competitive adsorption is caused by the dispersion forces whose interaction is established by Lennard-Jones and the electrostatic forces of gravity and repulsion, the mechanism of which is described by Van der Waals [8]. The competitive diffusion process involves two types of mass transfer: diffusion in the macropores (intercrystallite space) and diffusion in the micropores of crystallites (intracrystallite space).

2. During the evolution of the system towards equilibrium there is a concentration gradient in the macropores and in the micropores.

3. Competitive adsorption and diffusion occurs in active adsorbent centers, distributed throughout the inner surface of the nanoporous [7, 8]. All crystallites are spherical and have the same radius R , the crystallite bed is uniformly packed.

Taking into account the above, we have developed a nonlinear competitive adsorption model in the form of these hypotheses:

$$\frac{\partial C_j(t, Z)}{\partial t} = \frac{D_{inter_j}}{l^2} \frac{\partial^2 C_j}{\partial Z^2} - e_{inter_j} \frac{D_{intra_j}}{R^2} \left(\frac{\partial Q_j}{\partial X} \right)_{X=1} \quad (1)$$

$$\frac{\partial Q_j(t, X, Z)}{\partial t} = \frac{D_{intra_j}}{R^2} \left(\frac{\partial^2 Q_j}{\partial X^2} + \frac{2}{X} \frac{\partial Q_j}{\partial X} \right) \quad (2)$$

with initial conditions:

$$C_j(t, Z)|_{t=0} = 0, \quad Q_j(t, X, Z)|_{t=0} = 0, \quad X \in (0, 1), \quad j = \overline{1, 3} \quad (3)$$

and boundary conditions for coordinate X of the crystallite (particle):

$$\frac{\partial}{\partial X} Q_j(t, X, Z)|_{X=0} = 0. \quad (4)$$

In the expressions (1),(2) the function

$$Q_j(t, X = 1, Z)|_{X=1} = K_j C_j(t, Z) [1 + K_1 C_1(t, Z) + K_2 C_2(t, Z) + K_3 C_3(t, Z)]^{-1} \quad (5)$$

is the Langmuir competitive adsorption equilibria with boundary conditions on coordinate Z :

$$C_j(t, Z)|_{Z=1} = C_j^{in}, \quad \frac{\partial}{\partial Z} C_j(t, Z)|_{Z=0} = 0. \quad (6)$$

Here $c_j, q_j, j = \overline{1, 3}$ - is the current concentrations of the diffusion adsorbent components in the gas phase and micropores of the adsorbent particles, $c_{\infty j}, q_{\infty j}$ - is the corresponding equilibrium concentrations of the adsorbent components in the gas and adsorbed phase, $K_j = q_{\infty j}/c_{\infty j}$ - is the adsorption constant of the j -th component of the adsorbent, $K_j = 1/K_j, j = \overline{1, 3}$, ε_{inter} - the macroporosity of the media,

$$\varepsilon_{inter} = \frac{\varepsilon_{inter} c_j}{\varepsilon_{inter} c_j + (1 - \varepsilon_{inter}) q_j} \approx \frac{\varepsilon_{inter}}{(1 - \varepsilon_{inter}) K_j}.$$

Next, we perform the decomposition of a nonlinear system (1)-(5). Nonlinear function of the competitive Langmuir adsorption equilibrium is as follows [10]:

$$\varphi_j(C_1, C_2, C_3) = \frac{C_j(t, Z)}{1 + K_1 C_1(t, Z) + K_2 C_2(t, Z) + K_3 C_3(t, Z)}, \quad j = \overline{1, 3}. \quad (7)$$

We decompose into a series of Maclaurin at the point of zero concentrations for diffusion components of the adsorbate [13]:

$$\begin{aligned} \varphi_j^0(C_1, C_2, C_3) &= \varphi_j^0 + \left(\frac{\partial^2 \varphi_j^0}{\partial C_1 \partial C_2} C_1 C_2 + \frac{\partial^2 \varphi_j^0}{\partial C_1 \partial C_3} C_1 C_3 + \frac{\partial^2 \varphi_j^0}{\partial C_2 \partial C_3} C_2 C_3 \right) + \\ &+ \left(\frac{\partial \varphi_j^0}{\partial C_1} C_1 + \frac{\partial \varphi_j^0}{\partial C_2} C_2 + \frac{\partial \varphi_j^0}{\partial C_3} C_3 \right) + \frac{1}{2} \left(\frac{\partial^2 \varphi_j^0}{\partial C_1^2} C_1^2 + \frac{\partial^2 \varphi_j^0}{\partial C_2^2} C_2^2 + \frac{\partial^2 \varphi_j^0}{\partial C_3^2} C_3^2 \right) + \dots \end{aligned} \quad (8)$$

where $\varphi_j^0 = \varphi_j(0, 0, 0), j = \overline{1, 3}$. As a result, we obtain the following decompositions for (5) of the second order of accuracy:

$$\begin{aligned} Q_1(t, X=1, Z)|_{X=1} &= K_1 (C_1 - K_1 C_1^2 - K_2 C_1 C_2 - K_3 C_1 C_3), \\ Q_2(t, X=1, Z)|_{X=1} &= K_2 (C_2 - K_2 C_2^2 - K_1 C_1 C_2 - K_3 C_2 C_3), \\ Q_3(t, X=1, Z)|_{X=1} &= K_3 (C_3 - K_3 C_3^2 - K_1 C_1 C_3 - K_2 C_2 C_3). \end{aligned} \quad (9)$$

Assuming that $K_1 = \max\{K_j, K_j < 1\}_{j=\overline{1,3}}$, where $\varepsilon = K_1^2 \ll 1$ (small parameter), the problem (1)-(6), taking into account the approximated kinetic equations of phase transformation (9) containing a small parameter ε , is a boundary problem for a nonlinear system of partial differential equations. The solution of problem (1)-(5) will be sought with the help of asymptotic expansions in the small parameter ε in the form of the following series [13]:

$$C_j(t, Z) = C_{j0}(t, Z) + \varepsilon C_{j1}(t, Z) + \varepsilon^2 C_{j2}(t, Z) + \dots \quad (10)$$

$$Q_j(t, X, Z) = Q_{j0}(t, X, Z) + \varepsilon Q_{j1}(t, X, Z) + \varepsilon^2 Q_{j2}(t, X, Z) + \dots, \quad j = \overline{1, 3}. \quad (11)$$

As the result of substituting the asymptotic sum (10), (11) into equations (1)-(6) and replacing variables $N_{jm} = X \cdot Q_{jm}$, the initial nonlinear boundary problem (1)-(6) is parallelized into two types of linearized boundary problems.

The problem $A_{j_0}, j = \overline{1,3}$: to find a solution of partial differential equations system for the area $D = \{(t, X, Z) : t > 0, X \in (0, 1), Z \in (0, 1)\}$.

$$\frac{\partial C_{j_0}}{\partial t}(t, Z) = \frac{D_{inter_j}}{l^2} \frac{\partial^2 C_{j_0}}{\partial Z^2} - \epsilon_{inter_j} \frac{D_{intra_j}}{R^2} \left(\frac{\partial N_{j_0}}{\partial X} - N_{j_0} \right)_{X=1}, \quad (12)$$

$$\frac{\partial N_{j_0}}{\partial t}(t, X, Z) = \frac{D_{intra_j}}{R^2} \frac{\partial^2 N_{j_0}}{\partial X^2} \quad (13)$$

with initial conditions:

$$C_{j_0}(t, Z)|_{t=0} = 0; N_{j_0}(t, X, Z)|_{t=0} = 0; X \in (0, 1), j = \overline{1,3} \quad (14)$$

and boundary conditions for coordinate X of the crystallite:

$$N_{j_0}(t, X, Z)|_{X=0} = 0; N_{j_0}(t, X, Z)|_{X=1} = K_j C_{j_0}(t, Z), j = \overline{1,3} \quad (15)$$

The boundary conditions for coordinate Z are as follows:

$$C_{j_0}(t, Z)|_{Z=1} = 1; \frac{\partial}{\partial Z} C_{j_0}(t, Z)|_{Z=0} = 0. \quad (16)$$

The problem $A_m, m = \overline{1, \infty}$: to find in the area D a limited solution for the system of equations:

$$\frac{\partial C_{j_m}}{\partial t}(t, Z) = \frac{D_{inter_j}}{l^2} \frac{\partial^2 C_{j_m}}{\partial Z^2} - \epsilon_{inter_j} \frac{D_{intra_j}}{R^2} \left(\frac{\partial N_{j_m}}{\partial X} - N_{j_m} \right)_{X=1}, \quad (17)$$

$$\frac{\partial N_{j_m}}{\partial t}(t, X, Z) = \frac{D_{intra_j}}{R^2} \frac{\partial^2 N_{j_m}}{\partial X^2} \quad (18)$$

with zero initial conditions:

$$C_{j_m}(t, Z)|_{t=0} = 0; N_{j_m}(t, X, Z)|_{t=0} = 0, j = \overline{1,3} \quad (19)$$

boundary conditions for coordinate X of the crystallite:

$$N_{j_m}(t, X, Z)|_{X=0} = 0; N_{j_m}(t, X, Z)|_{X=1} = K_j C_{j_m}(t, Z) - F_{j_m}(t, Z), j = \overline{1,3} \quad (20)$$

$$F_{j_m}(t, Z) = \sum_{s=0}^{m-1} \sum_{k=1}^3 \frac{K_s K_k}{K_s^2} C_{j_s}(t, Z) C_{k_{m-1-s}}(t, Z), j = \overline{1,3}. \quad (21)$$

The boundary conditions for coordinate Z are as follows:

$$C_{j_m}(t, Z)|_{Z=1} = 0; \frac{\partial}{\partial Z} C_{j_m}(t, Z)|_{Z=0} = 0. \quad (22)$$

The construction methodology of an analytical solution. The Heaviside operating method [13] is used to find solutions C_{j_m} and Q_{j_m} of linearized system of problems (1)-(6). Assuming that the required functions C_{j_m} and Q_{j_m} ($N_{j_m} = X \cdot Q_{j_m}$) are Laplace originals, in Laplace images we obtain the next problems.

The problem A_0^* :

$$\frac{\partial^2 C_{j_0}^*}{\partial Z^2} = \frac{l^2}{D_{\text{inter}_j}} p C_{j_0}^* + \frac{3}{e_{\text{inter}_j}} \frac{l^2}{R^2} \frac{D_{\text{intra}_j}}{D_{\text{inter}_j}} \left(\frac{\partial N_{j_0}^*}{\partial X} - N_{j_0}^* \right)_{X=1}, \quad (23)$$

$$\frac{\partial^2 N_{j_m}^*}{\partial X^2} - \frac{R^2}{D_{\text{intra}_j}} p N_{j_m}^* = 0 \quad (24)$$

with boundary conditions for coordinate X of the crystallite:

$$N_{j_0}^*(p, X, Z)|_{X=0} = 0; N_{j_0}^*(p, X, Z)|_{X=1} = K_j C_{j_0}^*(p, Z), j = \overline{1, 3}. \quad (25)$$

The boundary conditions for coordinate Z are as follows:

$$C_{j_0}^*(p, Z)|_{Z=1} = C_{j_1}^*/p, \frac{\partial}{\partial Z} C_{j_0}^*(p, Z)|_{Z=0} = 0. \quad (26)$$

The problem $A_{j,m}^*$, $m = \overline{1, \infty}$:

$$\frac{\partial^2 C_{j_m}^*}{\partial Z^2} = \frac{l^2}{D_{\text{inter}_j}} p C_{j_m}^* + \frac{3}{e_{\text{inter}_j}} \frac{l^2}{R^2} \frac{D_{\text{intra}_j}}{D_{\text{inter}_j}} \left(\frac{\partial N_{j_m}^*}{\partial X} - N_{j_m}^* \right)_{X=1}, \quad (27)$$

$$\frac{\partial^2 N_{j_m}^*}{\partial X^2} - \frac{R^2}{D_{\text{intra}_j}} p N_{j_m}^* = 0. \quad (28)$$

The boundary conditions for coordinates X and Z are as follows:

$$N_{j_m}^*(p, X, Z)|_{X=0} = 0; N_{j_m}^*(p, X, Z)|_{X=1} = K_j C_{j_m}^*(p, Z) - F_{j_m}^*(p, Z), j = \overline{1, 3}. \quad (29)$$

$$C_{j_m}^*(p, Z)|_{Z=1} = 0; \frac{\partial}{\partial Z} C_{j_m}^*(p, Z)|_{Z=0} = 0. \quad (30)$$

Here $C_{j_m}^*(p, Z) = \int_0^\infty C_{j_m}(t, Z) e^{-pt} dt$, $N_{j_m}^*(p, X, Z) = \int_0^\infty N_{j_m}(t, X, Z) e^{-pt} dt$, p is the imaginary-significant parameter of Laplace transform.

The solution of the problem A_0^* . The solution of boundary problem (22)-(25) is:

$$N_{j_0}^*(p, X, Z) = K_j C_{j_0}^*(p, Z) \operatorname{sh} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} X \right) / \operatorname{sh} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right), j = \overline{1, 3}. \quad (31)$$

We calculate

$$\left(\frac{\partial N_{j_0}^*(p, X, Z)}{\partial X} \right)_{X=1} = R \sqrt{\frac{p}{D_{\text{intra}_j}}} \operatorname{cth} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right) K_j C_{j_0}^*(p, Z) \quad (32)$$

than, substituting (30) and (31) into equation (26) we obtain:

$$\frac{\partial^2 C_{j_0}^*}{\partial Z^2} - \gamma_j^2(p) C_{j_0}^*(p, Z) = 0 \quad (33)$$

where

$$\gamma_j^2(p) = \Gamma_j K_j \left(\frac{e_{\text{inter}_j}}{3K_j} \frac{R^2}{D_{\text{intra}_j}} p + R \sqrt{\frac{p}{D_{\text{intra}_j}}} \operatorname{cth} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right) - 1 \right);$$

$$\Gamma_j = \frac{3}{e_{\text{inter}_j}} \frac{l^2}{R^2} \frac{D_{\text{intra}_j}}{D_{\text{inter}_j}}$$

The solution of equation (22) taking into account the boundary conditions (29) are as follows:

$$C_{j_0}^*(p, Z) = C_j^m \frac{1}{p} \frac{\text{ch}[\gamma_j(p) Z]}{\text{ch}[\gamma_j(p)]} = C_j^m \frac{1}{p} \frac{\cos[\gamma_j(p) Z]}{\cos[\gamma_j(p)]} \quad (34)$$

The roots of transcendental equations $\text{ch}[\gamma_j(p)] = \cos[\gamma_j(\beta)] = 0$ are determined from transcendental equations $\gamma_j(\beta) = (2k - 1)\pi/2$, $k = \overline{1, \infty}$ or:

$$\beta^j \text{ctg}(\beta^j) = \frac{e_{\text{inter}_j}}{3K_j} (\beta^j)^2 = 1 - \frac{1}{\Gamma_j K_j} \left(\frac{2k - 1}{2} \pi \right)^2, \quad k = \overline{1, \infty} \quad (35)$$

Using Heaviside's theorem on the decomposition of a rational complex expression into a series by the roots of the denominator and making a substitution $p = -D_{\text{intra}_j} \beta_j^2 / R^2$, we obtain the formula for returning the Laplace image to the original (33):

$$C_{j_0}(t, Z) = C_j^m \left(1 + \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{\text{ch}[\gamma_j(p) Z] \exp(p_{ks} t)}{p_{ks} \frac{d}{dp} \text{ch}[\gamma_j(p)]|_{p=p_{ks}}} \right), \quad p_{ks} = -\frac{D_{\text{intra}_j} (\beta_{ks}^j)^2}{R^2} \quad (36)$$

where β_{ks}^j - the different positive roots of transcendental equations (34).

After simplifications we obtain:

$$C_{j_0}(t, Z) = C_j^m \left(1 + 2 \left(\frac{R}{l} \right)^2 \frac{D_{\text{inter}_j}}{D_{\text{intra}_j}} \times \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{(2k - 1) \pi \cos \left(\frac{2k - 1}{2} \pi Z \right) \exp \left(-\frac{D_{\text{intra}_j} (\beta_{ks}^j)^2 t}{R^2} \right)}{(-1)^s (\beta_{ks}^j)^2 \left[\frac{3K_j}{e_{\text{inter}_j}} \left(\frac{1}{\sin^2(\beta_{ks}^j)} - \frac{\text{ctg}(\beta_{ks}^j)}{\beta_{ks}^j} \right) - 2 \right]} \right) \quad (37)$$

Transforming the formula (30) to form we have:

$$N_{j_0}^*(p, X, Z) = C_{j_0}^*(p, Z) \frac{\sin(\beta X)}{\sin(\beta)} = \frac{C_j^m \text{ch}[\gamma_j(p) Z]}{p \text{ch}[\gamma_j(p)]} \frac{\sin(\beta X)}{\sin(\beta)}, \quad \beta = R \sqrt{\frac{p}{D_{\text{intra}_j}}}$$

and as a result of applying to it the Heaviside theorem on the expansion into a series, we obtain a formula for calculating the Laplace original of the function $N_{j_0}(t, X, Z)$:

$$N_{j_0}(t, X, Z) = 1 + \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{\sin(\beta X) \text{ch}[\gamma_{s1}(p_{kn}) Z] \exp(p_{kn} t)}{p_{kn} \sin(\beta) \frac{d}{dp} \text{ch}[\gamma_{s1}(p)]|_{p=p_{kn}}} \quad (38)$$

After transformation and going to the function $Q_{j_0}(t, X, Z)$ to describe the concentration distributions of the adsorbed components in the nanopores of the particles:

$$Q_{j_0}(t, Z) = C_j^m \left(1 + 2 \left(\frac{R}{l} \right)^2 \frac{D_{\text{inter}_j}}{D_{\text{intra}_j}} \times \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{(2k - 1) \pi \sin(\beta_{ks}^j X) \cos \left(\frac{2k - 1}{2} \pi Z \right) \exp \left(-\frac{D_{\text{intra}_j} (\beta_{ks}^j)^2 t}{R^2} \right)}{(-1)^s (\beta_{ks}^j)^2 X \sin(\beta_{ks}^j) \left[\frac{3K_j}{e_{\text{inter}_j}} \left(\frac{1}{\sin^2(\beta_{ks}^j)} - \frac{\text{ctg}(\beta_{ks}^j)}{\beta_{ks}^j} \right) + 2 \right]} \right) \quad (39)$$

The solution of the problem A_m . The solution of boundary problem (27), (28) is

$$N_{jm}^*(p, X, Z) = (K_j C_{jm}^*(p, Z) - F_{jm}^*(p, Z)) \frac{\text{sh} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} X \right)}{\text{sh} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right)}, j = \overline{1, 3} \quad (40)$$

From (39) at $X = 1$ we obtain:

$$\left(\frac{\partial N_{jm}^*(p, X, Z)}{\partial X} \right)_{X=1} = R \sqrt{\frac{p}{D_{\text{intra}_j}}} \text{cth} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right) (K_j C_{jm}^*(p, Z) - F_{jm}^*(p, Z)).$$

By substituting $N_{jm}^*|_{X=1}$ and $\frac{\partial N_{jm}^*}{\partial X}|_{X=1}$ in equation (38) we obtain:

$$\frac{d^2 C_{jm}^*}{dZ^2} - \gamma_j^2(p) C_{jm}^* = -\Phi_{jm}^*(p, Z) \quad (41)$$

here $\Phi_{jm}^*(p, Z) = \Gamma_j \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \text{cth} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right) - 1 \right) F_{jm}^*(p, Z)$, $m = \overline{1, \infty}$.

The solution of the inhomogeneous problem (40), (29) are as follows [15]:

$$C_{jm}^*(p, X, Z) = - \int_0^1 \mathcal{K}_j^*(p, Z, \xi) \Phi_{jm}^*(p, \xi) d\xi \quad (42)$$

where the fundamental function of Cauchy's influence $\mathcal{K}_j^*(p, Z, \xi)$, which is determined as:

$$\mathcal{K}_j^*(p, Z, \xi) = \begin{cases} \mathcal{K}_j^{*-} (p, Z, \xi) = d_1 \text{Ah} \gamma_j Z + e_1 \text{sh} \gamma_j Z, 0 < Z < \xi < 1 \\ \mathcal{K}_j^{*+} (p, Z, \xi) = d_2 \text{ch} \gamma_j Z + e_2 \text{sh} \gamma_j Z, 0 < \xi < Z < 1 \end{cases} \quad (43)$$

All the coefficients $d_s, e_s, s = \overline{1, 2}$ in expressions (44) are determined by the conditions [15]:

$$\begin{cases} \mathcal{K}_j^*(p, Z, \xi)|_{Z=\xi+0} - \mathcal{K}_j^*(p, Z, \xi)|_{Z=\xi-0} = 0 \\ \frac{d}{dZ} \mathcal{K}_j^*(p, Z, \xi)|_{Z=\xi+0} - \frac{d}{dZ} \mathcal{K}_j^*(p, Z, \xi)|_{Z=\xi-0} = 1 \end{cases} \quad (44)$$

and additional condition (boundary condition at $Z = 1$):

$$\mathcal{K}_j^{*+}|_{Z=1} = [d_2 \text{ch} \gamma_j Z + e_2 \text{sh} \gamma_j Z]_{Z=1} = 0. \quad (45)$$

Consistently applying the conditions conditions (44)-(46) as a result, the Cauchy function $\mathcal{K}_j^*(p, Z, \xi)$ is fully determined in the following form:

$$\mathcal{K}_j^*(p, Z, \xi) = -\frac{1}{\gamma_j} \begin{cases} \frac{\text{sh}(\gamma_j(1-\xi)) \cdot \text{ch}(\gamma_j Z)}{\text{ch}(\gamma_j)}, 0 < Z < \xi < 1 \\ \frac{\text{sh}(\gamma_j(1-Z)) \cdot \text{ch}(\gamma_j \xi)}{\text{ch}(\gamma_j)}, 0 < \xi < Z < 1 \end{cases} \quad (46)$$

Defining another function of influence:

$$\mathcal{H}_j^*(p, Z, \xi) = -\frac{R\sqrt{\frac{p}{D_{\text{intra}_j}}} \operatorname{cth}\left(R\sqrt{\frac{p}{D_{\text{intra}_j}}}\right)}{\gamma_j} \mathcal{K}_j^*(p, Z, \xi) \quad (47)$$

the solution (44) will take the following form :

$$C_{j_m}^*(p, X, Z) = -\Gamma_j \int_0^1 (\mathcal{H}_j^*(p, Z, \xi) - \mathcal{K}_j^*(p, Z, \xi)) F_{j_m}^*(p, \xi) d\xi. \quad (48)$$

The transition to originals. We carry out the transition to the originals $C_{j_m}(t, X, Z)$ using the formula:

$$C_{j_m}(t, X, Z) = -\Gamma_j \int_0^1 L^{-1} [\mathcal{H}_j^*(p, Z, \xi) - \mathcal{K}_j^*(p, Z, \xi)] * F_{j_m}(t, \xi) d\xi \quad (49)$$

where $L^{-1}[\dots]$ - is the designation of the Laplace inverse transform operator, $*$ is the im- convolution operator. In the final form, after calculating the originals, formula (47) will look like:

$$C_{j_m}(t, X, Z) = -\frac{3}{e_{\text{inter}_j}} \frac{l^2 D_{\text{intra}_j}}{\Pi^2 D_{\text{inter}_j}} * \int_0^1 \left(\int_0^1 (\mathcal{H}_j^-(t-\tau, Z, \xi) - \mathcal{K}_j^-(t-\tau, Z, \xi)) F_{j_m}^-(\tau, \xi) d\xi + \int_0^Z (\mathcal{H}_j^+(t-\tau, Z, \xi) - \mathcal{K}_j^+(t-\tau, Z, \xi)) F_{j_m}^+(\tau, \xi) d\xi \right) d\tau \quad (50)$$

where $\mathcal{H}_j^-(t-\tau, Z, \xi)$, $\mathcal{K}_j^-(t-\tau, Z, \xi)$, $\mathcal{H}_j^+(t-\tau, Z, \xi)$, $\mathcal{K}_j^+(t-\tau, Z, \xi)$ - are the components of influence functions (45), (47), the calculation algorithms of which are given below.

Applying to the components of the influence functions (45), (47) the Heaviside theorem on the development of a rational complex expression into a convergent series, we obtain:

$$L^{-1} \left[\frac{f_j^h(p)}{\gamma_j(p) \operatorname{sh}[\gamma_j(p)] \operatorname{ch}[\gamma_j(p)]} \right] = \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{f_j^h(\beta_{ks}^j) e^{\frac{D_{\text{intra}_j}}{R^2} (\beta_{ks}^j)^2 t}}{\omega_j^1(\beta_{ks}^j)} + \sum_{s_1=1}^{\infty} \frac{f_j^h(\mu_{s_1}^j) e^{\frac{D_{\text{intra}_j}}{R^2} (\mu_{s_1}^j)^2 t}}{\nu_j^2(\mu_{s_1}^j)} + \sum_{k_1=1}^{\infty} \frac{f_j^h(\eta_{k_1}^j) e^{\frac{D_{\text{intra}_j}}{R^2} (\eta_{k_1}^j)^2 t}}{\omega_j^2(\eta_{k_1}^j)} \quad (51)$$

where f_j^h , $h = \overline{1, 4}$ determine, respectively, the numerators of the components $\mathcal{H}_j^{\pm}(t-\tau, Z, \xi)$ - the influence functions (46).

Calculate the denominators in the expressions of the sums of each of the three terms of the right-hand side of formula (50):

$$\begin{aligned} \omega_j^1(\beta_{ks}^j) &\equiv \gamma_j(p) \operatorname{sh}\left(R\sqrt{\frac{p}{D_{\text{intra}_j}}}\right) \frac{d}{dp} \operatorname{ch}[\gamma_j(p)] \Big|_{p=\beta_{ks}^j} = \\ &= -\gamma_j(\beta_{ks}^j) \sin(\beta_{ks}^j) \frac{(-1)^k (\beta_{ks}^j)^2}{2k-1} \left[\frac{3K_j}{c_{\text{inter}_j}} \left(\frac{1}{\sin^2(\beta_{ks}^j)} - \frac{\operatorname{ctg}(\beta_{ks}^j)}{\beta_{ks}^j} \right) + 2 \right] \end{aligned} \quad (52)$$

where $\{\beta_{ks}^1\}$, $k, s = \overline{1, \infty}$ - set of roots of transcendental equation (34):

$$\begin{aligned} \nu_j^1(\mu_{s_1}^j) &\equiv \gamma_j(p) \operatorname{ch}[\gamma_j(p)] \frac{d}{dp} \operatorname{sh}\left(R\sqrt{\frac{p}{D_{\text{intra}_j}}}\right) \Big|_{p=-\frac{D_{\text{intra}_j}}{R^2}(\mu_{s_1}^j)^2} \equiv \\ &= \gamma_j(\mu_{s_1}^j) \cos[\gamma_j(\mu_{s_1}^j)] (-1)^{s_1} \frac{R^2}{2D_{\text{intra}_j} \mu_{s_1}^j} \Big|_{\mu_{s_1}^j = \pi s_1} \\ \omega_j^2(\eta_{k_1}^j) &\equiv \operatorname{sh}\left(R\sqrt{\frac{p}{D_{\text{intra}_j}}}\right) \operatorname{ch}[\gamma_j(p)] \frac{d}{dp} \gamma_j(p) \Big|_{p=-\frac{D_{\text{intra}_j}}{R^2}(\mu_{k_1}^j)^2} \equiv \\ &= \frac{1}{2R} \sqrt{\frac{3K_j}{e_{\text{inter}_j}} \frac{D_{\text{intra}_j}}{D_{\text{inter}_j}} \frac{\operatorname{ctg}(\eta_{k_1}^j)}{\eta_{k_1}^j} - \frac{1}{\sin^2 \eta_{k_1}^j} + \frac{2e_{\text{inter}_j}}{3}} \sin(\eta_{k_1}^j) \cos[\gamma_j(\eta_{k_1}^j)], \\ &\quad \sqrt{\frac{e_{\text{inter}_j}}{3K_j} + \frac{\operatorname{ctg}(\eta_{k_1}^j)}{\eta_{k_1}^j} - \frac{1}{(\eta_{k_1}^j)^2}} \end{aligned}$$

where $\{\eta_{k_2}^j\}_{k_2=1, \infty}^{j=1, 3}$ - set of roots of transcendental equation:

$$\frac{e_{\text{inter}_j}}{3K_j} (\mu^j)^2 - \mu^j \operatorname{ctg}(\mu^j) + 1 = 0, \quad \gamma_j^1(p) = 0, \quad R\sqrt{\frac{p}{D_{\text{intra}_j}}} = \mu^j$$

As a result, we obtain the following expressions of the originals:

$$\begin{aligned}
 \mathcal{H}_j^-(t, Z, \xi) &= L^{-1} [\mathcal{H}_j^{-*}(p, Z, \xi)] = \\
 &= \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{\beta_{ks}^j \cos(\beta_{ks}^j) \sin(\gamma_j(\beta_{ks}^j)(1-\xi)) \cdot \cos(\gamma_j(\beta_{ks}^j)Z) e^{\frac{D_{\text{intra}_j}(\beta_{ks}^j)^2 t}{R^2}}}{\omega_j^1(\beta_{ks}^j)} + \\
 &+ \sum_{s_1=1}^{\infty} \frac{\mu_{s_1}^j \cos(\mu_{s_1}^j) \sin(\gamma_j(\mu_{s_1}^j)(1-\xi)) \cdot \cos(\gamma_j(\mu_{s_1}^j)Z) e^{\frac{D_{\text{intra}_j}(\mu_{s_1}^j)^2 t}{R^2}}}{\nu_j^2(\mu_{s_1}^j)} \\
 &+ \sum_{k_1=1}^{\infty} \frac{\eta_{k_1}^j \cos(\eta_{k_1}^j) \sin(\gamma_j(\eta_{k_1}^j)(1-\xi)) \cdot \cos(\gamma_j(\eta_{k_1}^j)Z) e^{\frac{D_{\text{intra}_j}(\eta_{k_1}^j)^2 t}{R^2}}}{\omega_j^2(\eta_{k_1}^j)} \\
 \mathcal{H}_j^+(t, Z, \xi) &\equiv L^{-1} [\mathcal{H}_j^{+*}(p, Z, \xi)] = \\
 &= \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{\beta_{ks}^j \cos(\beta_{ks}^j) \sin(\gamma_j(\beta_{ks}^j)(1-Z)) \cdot \cos(\gamma_j(\beta_{ks}^j)\xi) e^{\frac{D_{\text{intra}_j}(\beta_{ks}^j)^2 t}{R^2}}}{\omega_j^1(\beta_{ks}^j)} + \\
 &+ \sum_{s_1=1}^{\infty} \frac{\mu_{s_1}^j \cos(\mu_{s_1}^j) \sin(\gamma_j(\mu_{s_1}^j)(1-Z)) \cdot \cos(\gamma_j(\mu_{s_1}^j)\xi) e^{\frac{D_{\text{intra}_j}(\mu_{s_1}^j)^2 t}{R^2}}}{\nu_j^2(\mu_{s_1}^j)} \\
 &+ \sum_{k_1=1}^{\infty} \frac{\eta_{k_1}^j \cos(\eta_{k_1}^j) \sin(\gamma_j(\eta_{k_1}^j)(1-Z)) \cdot \cos(\gamma_j(\eta_{k_1}^j)\xi) e^{\frac{D_{\text{intra}_j}(\eta_{k_1}^j)^2 t}{R^2}}}{\omega_j^2(\eta_{k_1}^j)}
 \end{aligned}$$

Next, we find the components $\mathcal{K}_j(t, Z, \xi)$. We have:

$$L^{-1} \left[\frac{g_j^h(p)}{\gamma_j(p) \operatorname{ch}[\gamma_j(p)]} \right] = \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{g_j^h(\beta_{ks}^j) e^{\frac{D_{\text{intra}_j}(\beta_{ks}^j)^2 t}{R^2}}}{\bar{\omega}_j^1(\beta_{ks}^j)} + \sum_{k_1=1}^{\infty} \frac{g_j^h(\eta_{k_1}^j) e^{\frac{D_{\text{intra}_j}(\eta_{k_1}^j)^2 t}{R^2}}}{\bar{\omega}_j^2(\eta_{k_1}^j)} \quad (53)$$

where $g_j^h(p)$ determine $\mathcal{K}_j^+(t - \tau, Z, \xi)$, $\mathcal{K}_j^-(t - \tau, Z, \xi)$ - the components of influence functions (45).

We calculate the denominators in the expressions of the sums of each of the two terms

of the right part of the formula (51):

$$\begin{aligned}
 \tilde{\omega}_j^1(\beta_{ks}^j) &\equiv \gamma_j(p) \frac{d}{dp} \operatorname{ch}[\gamma_j(p)] \Big|_{p=\beta_{ks}^j} = \\
 &= i\gamma_j(\beta_{ks}^j) \frac{(-1)^k \beta_{ks}^{j2}}{2k-1} \left[\frac{3K_j}{e_{\text{inter}_j}} \left(\frac{1}{\sin^2(\beta_{ks}^j)} - \frac{\operatorname{ctg}(\beta_{ks}^j)}{\beta_{ks}^j} \right) + 2 \right]; \\
 \tilde{\omega}_j^2(\eta_{k_1}^j) &= \operatorname{ch}[\gamma_j(p)] \frac{d}{dp} \gamma_j(p) \Big|_{p=-\frac{D_{\text{intra}_j}}{R^2} \eta_{k_1}^j} = \\
 &= \frac{l}{2R} \sqrt{\frac{3K_j D_{\text{intra}_j}}{e_{\text{inter}_j} D_{\text{inter}_j}}} \cdot \frac{\operatorname{ctg}(\eta_{k_1}^j) - \frac{1}{\sin^2 \eta_{k_1}^j} + \frac{2e_{\text{inter}_j}}{3}}{\sqrt{\frac{e_{\text{inter}_j}}{3K_j} + \frac{\operatorname{ctg}(\eta_{k_1}^j)}{\eta_{k_1}^j} - \frac{1}{(\eta_{k_1}^j)^2}}} = \cos[\gamma_j(\eta_{k_1}^j)]
 \end{aligned}$$

The expressions of originals are:

$$\begin{aligned}
 \mathcal{K}_j^-(t, Z, \xi) &= L^{-1}[\mathcal{K}_j^{-*}(p, Z, \xi)] = \\
 &= - \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{\sin(\gamma_j(\beta_{ks}^j)(1-\xi)) \cdot \cos(\gamma_j(\beta_{ks}^j)Z) e^{-\frac{D_{\text{intra}_j}(\beta_{ks}^j)^2}{R^2}t}}{\tilde{\omega}_j^1(\beta_{ks}^j)} \\
 &= - \sum_{k_1=1}^{\infty} \frac{\sin(\gamma_j(\eta_{k_1}^j)(1-\xi)) \cdot \cos(\gamma_j(\eta_{k_1}^j)Z) e^{-\frac{D_{\text{intra}_j}(\eta_{k_1}^j)^2}{R^2}t}}{\tilde{\omega}_j^2(\eta_{k_1}^j)} \\
 \mathcal{K}_j^+(t, Z, \xi) &= L^{-1}[\mathcal{K}_j^{+*}(p, Z, \xi)] = \\
 &= - \sum_{s=1}^{\infty} \sum_{k=1}^{\infty} \frac{\sin(\gamma_j(\beta_{ks}^j)(1-Z)) \cdot \cos(\gamma_j(\beta_{ks}^j)\xi) e^{-\frac{D_{\text{intra}_j}(\beta_{ks}^j)^2}{R^2}t}}{\tilde{\omega}_j^1(\beta_{ks}^j)} \\
 &= - \sum_{k_1=1}^{\infty} \frac{\sin(\gamma_j(\eta_{k_1}^j)(1-Z)) \cdot \cos(\gamma_j(\eta_{k_1}^j)\xi) e^{-\frac{D_{\text{intra}_j}(\eta_{k_1}^j)^2}{R^2}t}}{\tilde{\omega}_j^2(\eta_{k_1}^j)}
 \end{aligned}$$

$N_{j_m}(t, X, Z)$ is calculated on basis on the formula (51). We can obtain:

$$L^{-1}[N_{j_m}^*(p, X, Z)] = L^{-1}[K_j C_{j_m}^*(p, Z) - F_{j_m}^*(p, Z)] * L^{-1} \left[\frac{\operatorname{sh} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} X \right)}{\operatorname{sh} \left(R \sqrt{\frac{p}{D_{\text{intra}_j}}} \right)} \right] \quad (54)$$

We calculate the original of function $\Psi_{j_m}^*(p, X) = sh \left(R \sqrt{\frac{p}{D_{intra_j}}} X \right) / sh \left(R \sqrt{\frac{p}{D_{intra_j}}} \right)$:

$$L^{-1} [\Psi_{j_m}^*(p, X)] = \sum_{k_2=0}^{\infty} \frac{sh \left(R \sqrt{\frac{p}{D_{intra_j}}} X \right)}{\frac{d}{dp} \left(sh \left(R \sqrt{\frac{p}{D_{intra_j}}} \right) \right)_{p=p_{k_2}^j}} \exp(p_{k_2}^j t) =$$

$$- 2 \sum_{k_2=0}^{\infty} \frac{D_{intra_j}}{R^2} \frac{k_2 \pi \cdot \sin(k_2 \pi X)}{(-1)^{k_2+1}} \exp \left(- \frac{D_{intra_j}}{R^2} k_2^2 \pi^2 t \right)$$
(55)

Here $p_{k_2}^j = \frac{D_{intra_j}}{R^2} \pi^2 k_2^2$, $k_2 = \overline{0, \infty}$ are the roots of the equation $sh \left(R \sqrt{\frac{p}{D_{intra_j}}} \right) = 0$.

Applying the formula (50) in (49) one calculus the original $N_{j_m}(t, X, Z)$:

$$N_{j_m}(t, X, Z) = L^{-1} [(K_j C_{j_m}^*(p, Z) - F_{j_m}^*(p, Z)) * \Psi_j(p, Z)] -$$

$$= \int_0^t (K_j C_{j_m}(t - \tau, Z) - F_{j_m}(t - \tau, Z)) \times$$

$$\times \left(2 \frac{D_{intra_j}}{R^2} \sum_{k_2=0}^{\infty} \frac{\pi k_2 \cdot \sin(k_2 \pi X)}{(-1)^{k_2+1}} \exp \left(- \frac{D_{intra_j}}{R^2} k_2^2 \pi^2 t \right) \right) d\tau$$
(56)

Returning to the functions Q_{j_m} , we obtain:

$$Q_{j_m}(t, X, Z) = \int_0^t \left(K_j C_{j_m}(t - \tau, Z) - \sum_{s=0}^{m-1} \sum_{k=1}^3 \frac{K K_k}{K_F^k} C_k(t - \tau, Z) C_{k-1, \dots, 1}(t - \tau, Z) \right) \times$$

$$\times \left(2 \frac{D_{intra_j}}{R^2} \sum_{k_2=0}^{\infty} \frac{\pi k_2 \cdot \sin(k_2 \pi X)}{(-1)^{k_2+1}} \exp \left(- \frac{D_{intra_j}}{R^2} k_2^2 \pi^2 t \right) \right) d\tau, j = \overline{1, 3}.$$
(57)

The following theorem holds.

Theorem 1. *If the given and unknown functions of the boundary-value problems (23)-(23) and are Laplace pre-images with relative to the time variable t and the unique solvability conditions of boundary-value problems A_0 and A_m of the Laplace images are satisfied, then the solutions to boundary-value problems A_0 and A_m , exist, are unique, and are determined according to formulas (37),(39) and (51),(57), which constitute the solution of initial nonlinear boundary-value problem (1)-(6).*

2 SIMULATION AND DISCUSSION

The purpose of computer modeling was to study the capabilities of the model proposed for further use in technologies for cleaning carbon emissions into the atmosphere by energy and

Figure 1: Breakthrough curves ($ci(t)/c0$, $i = 1,2,3$) for three-component adsorption for a mixture of methane (CH_4 , $i = 1$), ethane (C_2H_6 , $i = 2$), propane (C_3H_8 , $i = 3$) taken from different mass ratios in the input mixture: (80%, 15%, 5%) (a), (35%, 35%, 30%) (b)

Figure 2: Desorption curves of gases in zeolite nanopores ($Q_i(t)/c0$, $i = 1,2,3$) for a mixture of methane (CH_4 , $i = 1$), ethane (C_2H_6 , $i = 2$), propane (C_3H_8 , $i = 3$) taken in different mass ratios in the input mixture: (80%, 15%, 5%) (a), (35%, 35%, 30%) (b)

transport facilities (propane, CO_2 and other combustion products). This is one of the key ways to solve the problem of global warming and create a safe energy strategy [2]. Propane was chosen as an adsorbent, the volume of which covers about 30% of the total gas flow leaving the car's engine. Using the developed mathematical theory and technology oriented to parallel multicore computer calculations, the modeling and calculation of concentration dependencies of three-component adsorption and desorption curves in nanoporous catalytic layers are carried out. Computational experiments were performed for the experimental sample [7, 10, 12].

3 CONCLUSIONS

High-performance methods and computational technologies for modeling non-isothermal gas adsorption in nanoporous solid for three-component adsorption equilibrium were developed. On their basis, new nonlinear mathematical models were constructed, including the balance equations of adsorption/desorption taking into account the interaction of intracrystallite space (micro flows) and intercrystallite space (macro flows). Implemented effective schemes for parallelizing and linearizing nonlinear models on basis of the decomposition of the equilibrium function into a series at the point of temperature of the phase transition as a small parameter. High-speed analytical and numerical solutions of mathematical models using the Heaviside operational method, their algorithmic and software implementation is implemented, which ensures efficient parallelization of computational processes for multicore computers and increased computational speed is constructed.

REFERENCES

- [Unused: 1] Unger N., Bond T.C., Wang J.S., Koch D.M., Menon S., Shindell D.T., Bauer S. *Attribution of climate forcing to economic sectors*. Proc. Natl. Acad. Sci. 2010, **107** (8), 3382–3389. doi:10.1073/pnas.0906548107
- [Unused: 2] Puertolas B., Navarro M.V., Lopez J.M., Murillo R., Mastral A.M., Garcia T. *Modelling the heat and mass transfers of propane onto a ZSM-5 zeolite*. Separation and Purification Technology 2012, **107** (2), 126–136. doi:10.1016/j.seppur.2011.10.036
- [Unused: 3] Krisnita R., Van Baten J.M. *Investigating the Non-idealities in Adsorption of CO₂-bearing Mixtures in Cation-exchanged Zeolites*. Separation and Purification Technology 2018, **206** (11), 208–217. doi:j.seppur.2018.06.009
- [Unused: 4] Santander J., Conner W.C., Jobic H., Auerbach S. *Simulating Microwave-Heated Open-Systems: Tuning Competitive Sorption in Zeolites*. J. Phys. Chem. B 2009, **113** (42), 13776–13781. doi:10.1021/jp902946g
- [Unused: 5] Hammond K.D., Conner W.C. *Analysis of Catalyst Surface Structure by Physical Sorption*. Advances in Catalysis. Springer-Verlag, Berlin, 2013.
- [Unused: 6] Karger J., Ruthven D., Theodorou D. *Diffusion in Nanoporous Materials*. John Wiley and Sons, Hoboken, 2012.
- [Unused: 7] Krishna R. *Thermodynamically Consistent Methodology for Estimation of Diffusivities of Mixtures of Guest Molecules in Microporous Materials*. ACS Omega 2019, **4** (8), 13520–13529. doi:10.1021/acsomega.9b01873
- [Unused: 8] Leclerc S., Petryk M., Canet D., Fraissard J. *Competitive Diffusion of Gases in a Zeolite Using Proton NMR and Slice Selection Procedure*. Catalysis Today 2012, **187** (1), 104–107. doi:10.1016/j.cattod.2011.09.007
- [Unused: 9] M. Petryk, M. Ivanchov, S. Leclerc, D. Canet and J. Fraissard. *Competitive Adsorption and Diffusion of Gases in a Microporous Solid*. IntechOpen Limited, London, 1973. doi: 10.5772/intechopen.88138
- [Unused: 10] Petryk M., Leclerc S., D. Canet, Sergienko I.V., Deineka V.S., Fraissard J. *The Competitive Diffusion of Gases in a zeolite bed: NMR and Slice Procedure, Modelling and Identification of Parameters* The Journal of Physical Chemistry C 2015, **119** (47), 26519–26525. doi:10.1021/acs.jpcc.5b07974
- [Unused: 11] Petryk, M.R., Boyko, I.V., Khimich, O.M., Petryk, M.M. *High-Performance Supercomputer Technologies of Simulation of Nanoporous Feedback Systems for Adsorption Gas Purification* Cybernetics and Systems Analysis 2020, **56** (5), 835–847. doi:10.1007/s10559-020-00304-y
- [Unused: 12] Petryk M., Khimitch A., Petryk M.M., Fraissard J. *Experimental and computer simulation studies of dehydration on microporous adsorbent of natural gas used as motor fuel* Fuel 2019, **239** (1), 1324–1330. doi:10.1016/j.fuel.2018.10.134
- [Unused: 13] Khimitch A., Petryk M., Mykhalyk D., Boyko I., Popov O., Sydoruk V. *Methods of the mathematical modeling and identification of complex processes and systems based on highly productive computing*. National Academy of Sciences of Ukraine. Glushkov Institute of Cybernetics, Kyiv, 2019.
- [Unused: 14] Lavrentiev M.A., Shabat B.V. *Methods of theory of functions of a complex variable*. Nauka, Moscow, 1973.
- [Unused: 15] V. V. Stepanov. *Course of the Differential Equations*. GIFML, Moscow, 1958.

Received 20.12.2020

Петрик М.Р., Бойко І.В., Шинкарик М.І., Петрик О.Ю. *Нелінійна модель конкурентивної дифузії для n-компонентного адсорбату з використанням рівноважної ізотерми Ленгмюра* // Буковинський матем. журнал — 2021. — Т.9, №1. — С. 10–24.

Наведені основи математичного моделювання неізотермічної трикомпонентної конкурентивної адсорбції газу в нанопористому середовищі з використанням рівноваги Ленгмюра. Запропоновано високоефективні аналітичні розв'язки для розширеної моделі адсорбції з використанням операційного методу Гевісайда та інтегрального перетворення Лапласа.

Наведено результати комп'ютерного моделювання на основі високшвидкісних обчислень на багатоядерних комп'ютерах.